


**AMENDMENTS TO THE CLAIMS**

This listing of claims will replace all prior versions, and listings, of claims in the application.

***Listing of Claims:***

1. (**Currently Amended**) A process for producing an N,N',N''-trisubstituted isocyanuric acid, comprising the step of:


heating an N-substituted carbamic acid derivative represented by following Formula (1):



wherein R is a hydroxyl-protecting group; and Z is a group represented by following Formula (2) or (3):



(2)




(3)

wherein R' is a hydrocarbon group or a heterocyclic group having a carbon atom at the bonding site with the adjacent oxygen atom,

wherein the heating step is carried out at a temperature in a range of 95°C to 145°C  
[[where]] when Z is the group represented by Formula (3),

to thereby form an N,N',N''-trisubstituted isocyanuric acid represented by following Formula (4):




wherein R has the same meaning as defined above.

2. **(Original)** A process for producing an N,N',N''-trisubstituted isocyanuric acid, comprising the step of heating an O-substituted hydroxylamine represented by following Formula (C):




wherein R is a hydroxyl-protecting group, or a salt thereof with (I) a compound represented by following Formula (A) or a compound represented by following Formula (B):



wherein X is a halogen atom; and Y is NH<sub>2</sub> or OR' wherein R' is a hydrocarbon group or a heterocyclic group having a carbon atom at the bonding site with the adjacent oxygen atom, or with (II) dimethyl carbonate, urea or phosgene, and a hydroxy compound represented by following Formula (D):




wherein R' has the same meaning as defined above to thereby form an N,N',N''-trisubstituted isocyanuric acid represented by following Formula (4):



wherein R has the same meaning as defined above.


3. **(Original)** A process for producing an N,N',N''-trisubstituted isocyanuric acid, comprising the steps of reacting a carbonyldiimidazole represented by following Formula (5):



with an O-substituted hydroxylamine represented by following Formula (6):



wherein R is a hydroxyl-protecting group, or a salt thereof, and further heating at a temperature in a range of 95°C to 145°C, to thereby form an N,N',N''-trisubstituted isocyanuric acid represented by following Formula (4):




wherein R has the same meaning as defined above.

4. **(Original)** The process for producing an N,N',N''-trisubstituted isocyanuric acid according to any one of claims 1 to 3, wherein the reaction is carried out in the presence of a base.

5. **(Currently Amended)** The process for producing an N,N',N''-trisubstituted isocyanuric acid according to claim 1, wherein R is an arylmethyl group which ~~may be~~ is optionally substituted.

6. **(Currently Amended)** The process for producing an N,N',N''-trisubstituted isocyanuric acid according to claim 1 or 2, wherein R' is an aromatic ~~eyelie~~ hydrocarbon group or aromatic heterocyclic group ~~which may be~~ wherein each group is optionally substituted.

7. **(Original)** A process for producing an N,N',N''-trisubstituted isocyanuric acid, comprising the step of purifying an N,N',N''-trisubstituted isocyanuric acid represented by following Formula (4):



wherein R is a hydroxyl-protecting group by at least one purification means selected from crystallization, repulping and washing, with the use of an alcohol-containing solvent.